

Clinical Policy: Alendronate (Binosto, Fosamax Plus D)

Reference Number: CP.PMN.88

Effective Date: 03.01.18

Last Review Date: 02.24

Line of Business: Commercial, HIM*, Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Alendronate sodium effervescent tablets (Binosto[®]) and alendronate/cholecalciferol (Fosamax Plus D[®]) are oral bisphosphonates.

**For Health Insurance Marketplace (HIM), Binosto is non-formulary and should not be approved using these criteria; refer to the formulary exception policy, HIM.PA.103.*

FDA Approved Indication(s)

Binosto and Fosamax Plus D are indicated for:

- Treatment of osteoporosis in postmenopausal women (PMO).
- Treatment to increase bone mass in men with osteoporosis.

Limitation(s) of use:

- Binosto and Fosamax Plus D: Optimal duration of use has not been determined. For patients at low-risk for fracture, consider drug discontinuation after 3 to 5 years of use.
- Fosamax Plus D alone should not be used to treat vitamin D deficiency.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Binosto and Fosamax Plus D are **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Osteoporosis (must meet all):

1. Diagnosis of PMO or male osteoporosis;
2. Age \geq 18 years or documentation of closed epiphyses on x-ray;
3. Failure of a 12-month trial of generic alendronate at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;
4. Dose does not exceed one of the following (a or b):
 - a. Binosto: 70 mg (1 tablet) per week;
 - b. Fosamax Plus D: 70 mg/5600 IU (1 tablet) per week.

Approval duration:

Medicaid/HIM – 12 months

Commercial – 12 months or duration of request, whichever is less

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Osteoporosis (must meet all):

1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
2. Member is responding positively to therapy;
3. If request is for a dose increase, new dose does not exceed one of the following (a or b):
 - a. Binosto: 70 mg (1 tablet) per week;
 - b. Fosamax Plus D: 70 mg/5600 IU (1 tablet) per week.

Approval duration:

Medicaid/HIM – 12 months

Commercial – 12 months or duration of request, whichever is less

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business:

CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or

2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid, or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

FDA: Food and Drug Administration

PMO: postmenopausal osteoporosis

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
alendronate (Fosamax [®])	<ul style="list-style-type: none"> • Treatment: PMO, male osteoporosis 10 mg PO QD or 70 mg PO once weekly • Prevention: PMO 5 mg PO QD or 35 mg PO once weekly 	40 mg/day 70 mg/week

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): abnormalities of the esophagus which delay esophageal emptying such as stricture or achalasia; inability to stand/sit upright for at least 30 minutes; hypocalcemia; hypersensitivity; increased risk of aspiration (Binosto only)
- Boxed warning(s): none reported

V. Dosage and Administration

Drug Name	Indication	Dosing Regimen	Maximum Dose
Alendronate effervescent (Binosto)	Treatment: PMO, male osteoporosis	70 mg PO once weekly	70 mg/week
Alendronate/cholecalciferol (Fosamax Plus D)		70 mg alendronate /2800 IU vitamin D3 or 70 mg	70 mg / 5600 IU/ week

Drug Name	Indication	Dosing Regimen	Maximum Dose
		alendronate /5600 IU vitamin D3 PO once weekly	

VI. Product Availability

Drug Name	Availability
Alendronate effervescent (Binosto)	Effervescent tablet: 70 mg
Alendronate/cholecalciferol (Fosamax Plus D)	Tablets: 70 mg/2800 IU, 70 mg/5600 IU

VII. References

1. Fosamax Plus D Prescribing Information. Jersey City, NJ: Organon LLC; August 2021. Available at: https://www.organon.com/product/usa/pi_circulars/f/fosamax/fosamax_plus_d_pi.pdf. Accessed October 19, 2023.
 2. Binosto Prescribing Information. Morristown, NJ: Ascend Therapeutics; October 2020. Available at: <https://www.binosto.com>. Accessed October 19, 2023.
 3. Clinical Pharmacology [database online]. Tampa, FL: Elsevier 2023. URL: www.clinicalkeys.com/pharmacology.
- Osteoporosis Diagnosis, Fracture Risk, and Treatment**
4. Shoback D, Rosen CJ, Black DM, et al. Pharmacological management of osteoporosis in postmenopausal women: an endocrine society guideline update. *J Clin Endocrinol Metab*; March 2020, 105(3): 587-594.
 5. Eastell R, Rosen CJ, Black DM, et al. Pharmacological management of osteoporosis in postmenopausal women: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab*; 2019, 104: 1595–1622.
 6. Camacho PM, Petak SM, Brinkley N et al. American Association of Clinical Endocrinologists/American College of Endocrinology clinical practice guidelines for the diagnosis and treatment of postmenopausal osteoporosis—2020 update. *Endocr Pract*. 2020;26(1):1-46.
 7. LeBogg MS, Greenspan SL, Insongna KL, et al. Clinician’s guide to prevention and treatment of osteoporosis. *Osteoporos Int*. 2022 Oct;33(10):2049-2102. doi: 10.1007/s00198-021-05900-y. Epub 2022 Apr 28. Erratum in: *Osteoporos Int*. 2022 Jul 28.
 8. Siris ES, Adler R, Bilezikian J, et al. The clinical diagnosis of osteoporosis: a position statement from the National Bone Health Alliance Working Group. *Osteoporos Int*. 2014 25:1439–1443. DOI 10.1007/s00198-014-2655-z.
 9. Hodsmann AB, Bauder DC, Dempster DW, et al. Parathyroid hormone and teriparatide for the treatment of osteoporosis: a review of the evidence and suggested guidelines for its use. *Endocr Rev*. 2005 Aug;26(5):688-703. Epub 2005 Mar 15.
- Male Osteoporosis**
10. Watts NB, Adler RA, Bilezikian JP, et al. Osteoporosis in men: an Endocrine Society clinical practice guidelines. *J Clin Endocrinol Metab* 2012;97(6):1802-1822.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
1Q 2020 annual review: added Medicaid line of business; age or closed epiphyses added; references reviewed and updated.	11.19.19	02.20
1Q 2021 annual review: no significant changes; references to HIM.PHAR.21 revised to HIM.PA.154; references reviewed and updated.	10.26.20	02.21
1Q 2022 annual review: no significant changes; references reviewed and updated.	09.13.21	02.22
Revised approval duration for Commercial line of business from length of benefit to 12 months or duration of request, whichever is less.	04.27.22	08.22
Template changes applied to other diagnoses/indications and continued therapy section.	10.10.22	
1Q 2023 annual review: no significant changes; references reviewed and updated.	11.01.22	02.23
1Q 2024 annual review: no significant changes; clarified failure of a “generic” alendronate is preferred; references reviewed and updated.	10.19.23	02.24

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan

retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.